

Dosing Charts

Note: 0.8 ml = 1 dropper

Acetaminophen (Tylenol) reduces fever and relieves pain. Typical uses include fevers, immunizations, teething, earaches, bruises, and headaches. Please notify us prior to using Tylenol for babies under 2 months old. Acetaminophen is available in the following forms:

• Infant Concentrated drops (80 mg/0.8 ml)

• Infant Oral Suspension (160 mg/5 ml) Note: 5 ml = 1 tsp

• Children's liquid (160 mg/5 ml)

• Chewable tablets (80 mg)

• Junior strength tablets (160 mg)

• Rectal suppository (80 mg, 120 mg, 325 mg) - can be used if the child is vomiting or very resistant to taking medicine by mouth. The suppositories can be cut-up to get the proper dose.

Acetaminophen (Tylenol) Dosing Chart - May give dose every 4 – 6 hours:

| Weight | Tylenol Milligram Dosage | Tylenol Infant drops 80mg/0.8 ml | Tylenol Infant Oral Suspension 160mg/5ml | Tylenol Children's liquid 160mg/5ml | Tylenol Chewable 80mg each | Tylenol Junior 160mg each |
|-------------|--------------------------------|--|---|--|----------------------------------|------------------------------|
| 6 – 8 lbs | 40 mg | ½ dropper (0.4 ml) | ¼ tsp (1.25 ml) | ½ tsp (1.25 ml) | N/A | N/A |
| 9 – 11 lbs | 60 mg | 3/4 dropper (0.6 ml) | 1/3 tsp (1.875 ml) | 1/3 tsp (1.875 ml) | N/A | N/A |
| 12 – 17 lbs | 80 mg | 1 dropper (0.8 ml) | ½ tsp (2.5 ml) | ½ tsp (2.5 ml) | N/A | N/A |
| 18 – 23 lbs | 120 mg | 1 ½ dropper (1.2 ml) | ³ / ₄ tsp (3.75 ml) | ³ / ₄ tsp (3.75 ml) | N/A | N/A |
| 24 – 35 lbs | 160 mg | 2 droppers (1.6 ml) | 1 tsp (5 ml) | 1 tsp (5 ml) | 2 tablets | 1 tablet |
| 36 – 47 lbs | 240 mg | 3 droppers (2.4 ml) | 1 ½ tsp (7.5 ml) | 1 ½ tsp (7.5 ml) | 3 tablets | 1 ½ tablet |
| 48 – 59 lbs | 320 mg | N/A | 2 tsp (10 ml) | 2 tsp (10 ml) | 4 tablets | 2 tablets |
| 60 – 71 lbs | 400 mg | N/A | 2 ½ tsp (12.5 ml) | 2 ½ tsp (12.5 ml) | 5 tablets | 2 ½ tablets |
| 72 – 95 lbs | 500 mg | N/A | 3 tsp (15 ml) | 3 tsp (15 ml) | 6 tablets | 3 tablets |

Please remember to keep all medications out of your child's reach at all times



Dosing Charts

Ibuprofen (Motrin, Advil) reduces fever, relieves pain, and fights inflammation. Typical uses include fevers, immunizations, teething, earaches, bruises, headache, sunburns, and sore muscles. We usually recommend that ibuprofen not be used in babies under 6 months old. Ibuprofen is available in the following forms:

Infant drops (50mg/1.25ml)

• Children's liquid (100mg/5ml)

• Chewable tablets (50mg)

Junior strength tablets (100mg)

Note: 1.25ml= 1 dropper

Note: 5 ml = 1 tsp

Ibuprofen (Motrin/Advil) Dosing Chart - May give dose every 6-8 hours:

| Weight | Motrin Milligram Dosage | Motrin Children's liquid 100mg/5ml | Motrin Chewable 50mg each | Motrin Junior 100mg each |
|-------------|-------------------------------|--|---------------------------------|-----------------------------------|
| 12 – 17 lbs | 50 mg | ½ tsp (2.5 ml) | N/A | N/A |
| 18 – 23 lbs | 75 mg | ³ / ₄ tsp (3.75 ml) | N/A | N/A |
| 24 – 35 lbs | 100 mg | 1 tsp (5 ml) | 2 tablets | 1 tablet |
| 36 – 47 lbs | 150 mg | 1 ½ tsp (7.5 ml) | 3 tablets | 1 ½ tablet |
| 48 – 59 lbs | 200 mg | 2 tsp (10 ml) | 4 tablets | 2 tablets |

Please remember to keep all medications out of your child's reach at all times